

# GUIDANCE FRAMEWORK FOR BETTER AIR QUALITY IN ASIAN CITIES

Guidance Area 3: Health and other impacts





# **GUIDANCE FRAMEWORK FOR BETTER AIR QUALITY IN ASIAN CITIES**

Guidance Area 3: Health and other impacts

---

## ABOUT CLEAN AIR ASIA

[www.cleanairasia.org](http://www.cleanairasia.org)

Clean Air Asia is an international NGO established in 2001 as the premier air quality network for Asia by the Asian Development Bank, World Bank and USAID. Its mission is to promote better air quality and livable cities by translating knowledge to policies and actions that reduce air pollution and greenhouse gas emissions from transport, energy and other sectors.

Clean Air Asia became a UN-recognized partnership in 2007, its network spanning 250 organizations in 31 countries in Asia and worldwide, with 8 country networks: China, India, Indonesia, Nepal, Pakistan, Philippines, Sri Lanka and Vietnam. It is headquartered in Manila and has offices in Beijing and Delhi. Clean Air Asia leads efforts to enable Asia's more than 1000 cities to reduce both air pollution and CO<sub>2</sub> emissions, and thereby contribute to more livable and healthy cities with blue skies and low carbon footprint. Clean Air Asia helps to reduce emissions through policies, plans, programs, and concrete measures that cover air quality, transport and industrial emissions and energy use.

The Better Air Quality (BAQ) Conference is a flagship event of Clean Air Asia covering the key sectors of transport, energy and industry, with a particular emphasis on government policies and measures. Policy makers, practitioners and industry leaders meet at BAQ to network, innovate, learn and share experiences. The biennial event was first held in 2002 and attracts close to a thousand participants from Asia and the rest of the world.

©2016 Clean Air Asia. All rights reserved.  
Published by Clean Air Asia, 2016. "Guidance Framework for Better Air Quality in Asian Cities". Pasig City, Philippines.  
ISBN: 978-621-95434-0-8

Photo credits: iStock, Shutterstock, Design Muscle Inc., Clean Air Asia and Ms. Maria Katherina Patdu

This publication may be reproduced in whole or in part in any form for educational or non-profit purposes without special permission from the copyright holder, provided acknowledgment of the source is made. The Clean Air Asia Center would appreciate receiving a copy of any publication that uses this Clean Air Asia Center publication as a source. No use of this publication may be made for resale or for any other commercial purpose whatsoever, without prior permission in writing from the Clean Air Asia Center.

### Disclaimer

The views expressed in this publication are those of Clean Air Asia staff, consultants and management, and do not necessarily reflect the views of the Board of Trustees of the Clean Air Asia Center. The Clean Air Asia Center does not guarantee the accuracy of the data included in this publication and does not accept responsibility for consequence of their use.

### Contact

**Clean Air Asia Center**  
Unit 3505 Robinsons Equitable Tower  
ADB Avenue, Pasig City, 1605  
Philippines  
Tel +632 6311042  
Fax +63 2 6311390  
[center@cleanairasia.org](mailto:center@cleanairasia.org)

**Clean Air Asia China Office**  
11152, JianGuoMenWai  
Diplomatic Residence  
Compound, No.1 XiuShui Street,  
ChaoYang District, Beijing,  
China, 100600  
Tel/Fax:+86 10 8532 6972  
[china@cleanairasia.org](mailto:china@cleanairasia.org)

**Clean Air Asia India Office**  
1st Floor, Building No.4  
Thyagraj Nagar Market Lodhi Colony  
110003, India  
Tel +91 11 60120260  
Fax +91 11 43850032  
[india@cleanairasia.org](mailto:india@cleanairasia.org)

**Country Networks in China,  
India, Indonesia, Nepal,  
Pakistan, Philippines, Sri  
Lanka, Vietnam**

## CONTENTS

Preface	vi
Abbreviations	vii
4.1 Introduction	1
4.1.1 Objective	2
4.1.2 Importance of considering health and environmental impacts	2
4.1.2.1 Health impacts of air pollution	4
4.1.2.2 Environmental impacts of air pollution	6
4.1.3 Link of air pollution and climate change; co-benefits	8
4.1.4 Health impact assessment in clean air action plans	9
4.2 Stages of health and environmental impact assessment	10
4.3 Issues and challenges	14
4.4 Roadmap for health and environmental impact assessment	16
4.5 References	21

## PREFACE

Air pollution is now considered the world's largest environmental health risk. There have been a number of global efforts calling for air pollution actions in recent years. These global calls for action on air pollution strengthen regional and national initiatives and highlight the need to prioritize addressing this issue through a collaborative and integrated approach.

In 2006, the First Governmental Meeting on Urban Air Quality in Asia<sup>1</sup> recognized the need for guidance in implementing a Long Term Vision for Urban Air Quality in Asia, which describes the desired state of urban air quality management in Asian cities. During the Third Governmental Meeting, environment ministries from the region identified key challenges they are facing to improve urban air quality.

To set the way forward in achieving the vision for cleaner air, Clean Air Asia led the development of the Guidance Framework for Better Air Quality in Asian Cities (Guidance Framework) in 2014 to address the needs and challenges in the region. It aims to provide a recognized guidance on improving urban air quality and is organized around priority areas of concern in the region, which were translated into key guidance areas with roadmaps on how to progress in a step by step manner.

This voluntary, non-binding document consists of seven individually published chapters covering each of the Guidance Areas. Policy and decision makers in Asia, as well as other relevant stakeholders, can use one or a combination of the Guidance Framework chapters to develop local roadmaps or action plans depending on their priority areas of concern.

The Guidance Framework consists of seven main books with these titles:

- Introduction
- Guidance Area 1 - Ambient air quality standards and monitoring
- Guidance Area 2 - Emissions inventories and modeling
- Guidance Area 3 - Health and other impacts
- Guidance Area 4 - Air quality communication
- Guidance Area 5 - Clean air action plans
- Guidance Area 6 - Governance

These guidance areas come with an Information Sourcebook, which is a compilation of resources to support the implementation of Guidance Framework roadmaps. There is also an accompanying training course on Guidance Framework implementation, which is available online in the Clean Air Asia website and Integrated Programme for Better Air Quality (IBAQ Programme) website: [www.cleanairasia.org/ibaq](http://www.cleanairasia.org/ibaq)

The Guidance Framework was developed together with a team of international and regional experts and practitioners and has undergone an extensive review process through the Governmental Meetings and the involvement of external reviewers. The draft document was also shared in a number of international events, including the Asia Pacific Clean Air Partnership (APCAP) Joint Forum organized by UNEP ROAP in November 2015. The Guidance Framework was welcomed by participants from 24 countries in Asia and the Pacific, involving environment ministries, intergovernmental organizations, non-governmental organizations, and experts.

## ABBREVIATIONS

ABCs	Atmospheric Brown Clouds	MOEF (India)	Ministry of Environment, Forest, and Climate Change
AQM	Air Quality Management	NAAQS	National Ambient Air Quality Standards
BC	Black Carbon	NAMP	National Air Quality and Monitoring Program
CAAP	Clean Air Action Plan	NGO	Non-Governmental Organization
CAHA	Climate and Health Alliance	NH <sub>3</sub>	Ammonia
CPCB	Central Pollution Control Board	NOx	Nitrogen oxides
CFCs	Chlorofluorocarbons	NO <sub>2</sub>	Nitrogen dioxide
CH <sub>4</sub>	Methane	OECD	Organisation for Economic Co-operation and Development
CO	Carbon monoxide	O <sub>3</sub>	Ozone
CO <sub>2</sub>	Carbon dioxide	PFC	Perfluorinated Compounds
DPSEEA	Driving Force-Pressure-State-Exposure-Effect-Action	PM	Particulate Matter
ECHP	European Centre for Health Policy	PM <sub>10</sub>	Particulate Matter (≤ 10 micrometers in diameter)
EPD (Hong Kong)	Environmental Protection Department	PM <sub>2.5</sub>	Particulate Matter (≤ 2.5 micrometers in diameter)
GAPF	Global Atmospheric Pollution Forum	SEI	Stockholm Environment Institute
GBD	Global Burden of Disease	SLCPs	Short-Lived Climate Pollutants
GCEC	Global Commission on the Economy and Climate	SO <sub>2</sub>	Sulfur dioxide
GHG	Greenhouse Gas	UK	United Kingdom
HCFC	Hydrochlorofluorocarbons	UNEP	United Nations Environment Programme
HFC	Hydrofluorocarbons	USEPA	United States Environmental Protection Agency
HFE	Fluorinated ethers	WHO	World Health Organization
HIA	Health Impact Assessment	WMO	World Meteorological Organization
IARC	International Agency for Research on Cancer		
IPCC	International Panel on Climate Change		

1 Governmental Meetings on Urban Air Quality in Asia are biennial meetings organized by the United Nations Environment Programme Regional Office of Asia and the Pacific (UNEP ROAP) and Clean Air Asia that convene environment ministries with the aim to harmonize approaches across the region in tackling air pollution and related fields.



The exposure to ambient air pollution, particularly fine particulate matter is estimated by WHO to have caused 3.7 million premature deaths worldwide in 2012.

## CHAPTER 4

### GUIDANCE AREA 3: HEALTH AND OTHER IMPACTS

#### 4.1 Introduction

Ambient air pollution is a much more significant public health risk than previously assumed. This is one of the findings of two recent estimates on the global impact of air pollution on health—from the World Health Organization (WHO) in 2012 (WHO, 2014a) and from the Global Burden of Disease (GBD) project in 2010 (Lim et al., 2012).

The exposure to ambient air pollution, particularly fine particle or particulate matter (PM<sub>2.5</sub>) is estimated by WHO to have caused 3.7 million<sup>1</sup> premature deaths worldwide in 2012. Most of the burden of air pollution is borne by the middle- and low-income countries in the WHO Western Pacific (WHO, 2014b) and South-East Asia regions – an estimated 1.6 million and 0.94 million deaths, respectively.

Over 76 million years of healthy life are lost due to air pollution according to the GBD's 2010 findings (Lim et al., 2012). The GBD ranks air pollution as the third most important risk of premature death in the world, after dietary risks and high blood pressure.

<sup>1</sup> WHO's estimates' range is 3.188-4.291 million.

**To establish and/or strengthen national and local programs that monitor the health, environmental and economic impact of air pollution consistently.**

The GBD study shows that Asia gets the largest share of the ambient PM<sub>2.5</sub>-related deaths worldwide with an estimated 2.15 million deaths in 2010. Approximately 57 percent of these cases are in China and about 29 percent in India. This is also the case with premature deaths worldwide due to ground-level ozone (O<sub>3</sub>) exposure, with Asia accounting for 83 percent of such deaths or 126,508 cases in 2010 (Figure 4.1).

More recently, in 2013, a group of experts working on behalf of WHO's International Agency for Research on Cancer (IARC) classified ambient air pollution in general and PM<sub>2.5</sub> in particular as carcinogenic (IARC Group 1) to humans (WHO/IARC, 2013a, b, c). This classification is based on sufficient evidence of carcinogenicity in humans and experimental animals, and strong evidence on the mechanisms (Brauer et al., 2012; Loomis et al., 2013; 2014). As a carcinogen, particulate matter (PM) has no threshold for the onset of effects.




Figure 4.1 Regional distribution of premature deaths per year due to PM<sub>2.5</sub> and O<sub>3</sub>

Source: Lim et al., 2012

**4.1.1 Objective**

To establish and/or strengthen national and local programs that monitor the health, environmental, and economic impact of air pollution consistently.

pollutants have been classified as carcinogenic, in addition to causing a variety of respiratory and cardiovascular diseases. Impacts of air pollution on the environment include acidification of soil and freshwater systems, eutrophication of lakes and rivers, ozone layer depletion and global climate change.

**4.1.2 Importance of considering health and environmental impacts**

Ambient and household air pollution has many significant impacts on human health and the environment. Many air

Understanding and estimating the impacts of air pollution as well as its economic costs to society are integral in the overall air quality management (AQM) framework. Any AQM framework is based on the Driving force-Pressure-State-Exposure-Effect-Action (DPSEEA) framework depicted in Figure 4.2.

Analyzing the cost of air pollutant control measures *vis a vis* the benefits of avoided health and environmental impacts as a result of these measures is critical in improving the effectiveness of AQM. Without estimates on pollution damage costs, it is impossible to determine the cost

effectiveness of air pollution prevention, mitigation, and control strategies. Identifying efficient and effective technologies and policy tools, needed for target setting and management strategy, also becomes unworkable.




Figure 4.2 The DPSEEA framework<sup>2,3</sup>

Source: Adapted from WHO, 1995a

**Understanding and estimating the impacts of air pollution as well as its economic costs to society are integral in the overall air quality management framework.**

2 Absorbed dose: The amount of air pollutant absorbed, and therefore available to undergo metabolism, transport, storage, or elimination in the human body (WHO, 2000b).  
 3 Pathogen intake: The taking in of substances which can cause harm or disease (WHO, 2000a; Cyclopedic Medical Dictionary, 1997).



### 4.1.2.1 Health impacts of air pollution

The greatest impacts on human health tend to be focused on the effects of these common pollutants – PM<sub>2.5</sub>, O<sub>3</sub>, nitrogen dioxide (NO<sub>2</sub>), and sulfur dioxide (SO<sub>2</sub>) – as well as toxic air pollutants and heavy metals.

#### Particulate matter

Evidence suggests that there is a close correlation between exposure to high concentrations of small particles and increased mortality or morbidity, both in the short- and long-term (Hirota & Martin, 2013; United States Environmental Protection Agency [USEPA], 2009; Hogg et al., 2004; WHO, 2013a,b; Brook et al., 2010; Chuang et al., 2007; R ckerl et al., 2006). Fine particulate matter, after it is inhaled, penetrates deep into the lung and the air sacs. The particles damage cells in the airways and affect the lung; this is associated with the exacerbation of asthma and chronic obstructive pulmonary disease (Riva et al., 2011; USEPA, 2009; Risom et al., 2005). The fine particulates cause further damage as they impact the heart and circulatory system.

PM<sub>2.5</sub>-related deaths occur due to:

- cardiovascular disease, notably ischemic heart disease and stroke (WHO, 2013a,b);
- lung cancer (WHO, 2013a, b, c);
- chronic obstructive pulmonary disease (WHO, 2013a, b, c); and
- acute lower respiratory infection (WHO, 2013a, b; Roth et al., 2008; Lanata et al., 2004).

Non-fatal health impacts of exposure to PM<sub>2.5</sub> include non-fatal heart attacks, irregular heartbeat, aggravated asthma and decreased lung function, acute bronchitis, and increased respiratory symptoms. These can lead to absence from work and reduced school attendance, as well as increased hospital admissions and doctor’s visits.

A component of PM<sub>2.5</sub> that should also be mentioned is black carbon (BC), which refers to elemental carbon-containing particles that are emitted from incomplete combustion of fossil fuels and biomass (Long et al., 2013). According to WHO, there is sufficient evidence for an association between daily variations in BC concentrations and short-term changes in all-cause and cardiovascular mortality, as well as cardiopulmonary hospital admissions (Janssen et al., 2012). For long-term BC exposure, cohort studies have provided sufficient evidence of relationships to all-cause and cardiopulmonary mortality. Exposure-relationships from both short- and long-term studies are much stronger for BC than for PM<sub>2.5</sub> and PM<sub>10</sub> (Janssen et al. 2012). This statement does not mean that “any specific source, component, or size class of PM maybe excluded as a possible contributor to PM toxicity” (Hedley Index, 2013).

Given that ambient PM<sub>2.5</sub>-related deaths are largely concentrated in Asia, the risk of ambient PM ranks among the top health risks everywhere in Asia (Table 4.1).

Table 4.1 Risk factors ranked by attributable burden of disease, 2010

	Global	High-income Asia Pacific	East Asia	South East Asia	South Asia	Central Asia
High blood pressure	1	1	1	1	3	1
Tobacco smoking, including second-hand smoke	2	2	2	2	2	3
Alcohol use	3	3	6	6	8	2
Household air pollution from solid fuels	4	42	5	3	1	12
Diet low in fruits	5	5	3	4	5	5
High body-mass index	6	8	9	9	17	4
High fasting plasma glucose	7	7	8	5	7	7
Childhood underweight	8	39	38	13	4	25
Ambient particulate matter pollution	9	9	4	11	6	10
Physical inactivity and low physical activity	10	4	10	8	11	9

Source: Lim et al., 2012

#### Ground-level ozone

Ground-level ozone is less soluble in water. It is thus not scrubbed in the upper respiratory tract and reaches the lower respiratory tract where it dissolves in the fluid of the thin surface layer of lung cells. Free radicals and other oxidants in the fluid are assumed to react rapidly with the cell molecules and mediate the effects of O<sub>3</sub> exposure of the human lung. Ozone exposure is a major factor in asthma morbidity and mortality. Even short-term exposure can aggravate existing lung diseases and make the lungs more susceptible to infection.

Health effects of short-term exposure to O<sub>3</sub> include (USEPA, 2014a, 2013a; WHO 2013a, b):

- Increased all-cause mortality
- Increased cardiovascular mortality in adults younger than 75 years

- Increased hospital admission for heart diseases in adults older than 65 years
- Increased hospital admissions for respiratory diseases in adults older than 65 years
- Increased hospital admissions for chronic obstructive pulmonary disease
- Potentially increased hospital admissions for asthma
- Increased school absences

#### Nitrogen dioxide

Upon inhalation of NO<sub>2</sub>, the human respiratory tract can absorb between 70-90 percent of the gas, and even greater levels when breathing from the mouth – e.g., during exercise. Nitrogen dioxide is increasingly deposited in the lower respiratory tract and can remain in the lung for prolonged periods (WHO, 2006). Short-term exposure to NO<sub>2</sub> is enough to cause aggravations in people with asthma and other



lung diseases. Above safe levels, NO<sub>2</sub> exposure can cause inflammation of the airways. Some studies have shown associations between NO<sub>2</sub> long-term exposure and mortality. However, present evidence is not sufficient to conclude that effects on mortality can be attributed to exposure to NO<sub>2</sub> itself (Institute of Occupational Medicine, 2004).

Health effects of long-term exposure to NO<sub>2</sub>:

- Exacerbation of symptoms of bronchitis in asthmatic children (WHO, 2013b)
- Reduced lung function in children (WHO, 2013a; b)

#### Sulfur dioxide

The main anthropogenic source of SO<sub>2</sub> is the burning of sulfur-containing fossil fuels for domestic heating, power generation and motor vehicles. Being highly soluble in water, inhaled SO<sub>2</sub> is readily absorbed in the human respiratory tract, in particular, while breathing from the mouth. Inhaled SO<sub>2</sub> can cause inflammation of the respiratory tract (WHO, 2013a).

Epidemiological studies have linked SO<sub>2</sub> long-term exposure to:

- Causation of changes in lung function
- Exacerbation of existing heart diseases
- Increased number of asthma attacks

WHO, in its most recent factsheet (WHO, 2014), believes that present evidence is sufficient to conclude that effects on mortality can be attributed to exposure to SO<sub>2</sub>.

#### Toxic air pollutants and heavy metals

Examples of toxic air pollutants include benzene (in gasoline), benzo[a]pyrene (in tar and asphalt fumes, and diesel engine emissions), perchloroethylene (in dry cleaning chemicals), methylene chloride (in solvents and paint strippers), asbestos, toluene, arsenic, and metals such as zinc, cadmium, mercury, chromium, nickel, and lead compounds.

People exposed to toxic air pollutants due to poisonous smog at ample concentrations and durations may have increased chances of cancer, reproductive and birth defects, immunological damages, and retarded neurological development. Some persistent toxic air pollutants accumulate in body tissues because unlike organic pollutants, toxic metals do not decay.

Toxic air pollutants such as benzene, benzo[a]pyrene, nickel and arsenic, in particular, are classified as carcinogenic; no safe level of exposure has been derived by WHO for these metals. While ammonia (NH<sub>3</sub>) in itself is not classified as carcinogenic, it converts gaseous acids, especially sulfuric and nitric acids, to PM<sub>2.5</sub>, which is carcinogenic (Martin, 2008). In a recent publication, Paulot & Jacob (2014) suggested that elimination of NH<sub>3</sub> emissions would achieve a greater health benefit than the reduction of the PM<sub>2.5</sub> national standard from 15 to 12 µg/m<sup>3</sup>.

More detailed information about health impacts due to exposure to toxic air pollutants can be accessed at WHO (1995b) and USEPA (2007).

#### 4.1.2.2 Environmental impacts of air pollution

Air pollutants are causing changes in the ecosystem and are indirectly affecting wildlife, in addition to directly affecting animal populations that are exposed to harmful air pollutants (United Kingdom Website (UK), 2014b). Extensive experimental studies to assess the potential threat from O<sub>3</sub> to agriculture conducted in Europe and North America have demonstrated crop losses due to O<sub>3</sub> impacts amounting to billions of dollars (Avnery et al., 2011a). The impact of air pollution on cultural heritage materials is also a serious concern because it can lead to loss of important parts of a county's history and culture.

#### Acid rain

Acid rain containing nitric and sulfuric acids, causes damage to forests and the acidification of soils and water bodies, potentially disrupting the food chain. Acid deposition may have various effects on crops, including significant yield loss in many species. Acid rain also accelerates the erosion of buildings, statues, and sculptures. Further information can be accessed from the USEPA (2012b), Phinney et al. (2004), and UK (2014a).

#### Eutrophication

Dry and wet deposition of airborne nutrients such as nitrates and phosphates can stimulate algal blooms. In turn, these blooms can cause fish kills and loss of plant and animal life because of oxygen depletion (United States Geological Survey, 2014).

#### Haze

Haze is caused by the scattering and absorption of sunlight by airborne fine PM (USEPA, 2012c). Haze degrades visual range, obscuring the clarity, color, texture, and forms of objects in cities and scenic areas. Haze has potential impacts on air quality, climate, and the hydrological cycle (Ramanathan et al., 2001).

#### Atmospheric brown clouds

Atmospheric brown clouds (ABCs) primarily consist of particles and pollutant gases, such as nitrogen oxides (NOx), carbon monoxide (CO), SO<sub>2</sub>, NH<sub>3</sub>, and hundreds of organic gases and acids from indoor and outdoor air pollution due to biomass and fossil fuel burning (Ramanathan et al., 2008). Five ABC hotspots exist; three of them lie in East, South, and South-East Asia. The five ABC hotspots potentially have large impacts on regional and global climate, freshwater budget, agriculture, and health (Ramanathan et al., 2008; Ramanathan & Ramana, 2003).

#### Ozone layer depletion

Stratospheric ozone, particularly in the polar zones, is being gradually destroyed by chlorine- and bromine-containing substances referred to as ozone depleting substances. The Montreal Protocol on Substances that Deplete the Ozone Layer entered into force in 1989 to protect stratospheric ozone and, consequently, protect life from increased ultraviolet radiation at Earth's surface (United Nations Environment Programme [UNEP], 2011). These controls under the Protocol have proven effective for safeguarding public health<sup>4</sup>. Ultraviolet radiation can damage sensitive crops and reduce crop yields. Most ozone-depleting substances are also potent greenhouse gases, contributing to global warming. There is strong evidence available on the effect of stratospheric ozone changes on Earth's surface climate (UNEP, 2010; USEPA, 2011a).

#### Crop and forest damage

Ground-level O<sub>3</sub> can reduce agricultural crop and commercial forest yields, diminish growth and influence survival of tree seedlings, and increase plant susceptibility to disease, pests and other stresses. Wang & Mauzerall (2004) estimated a total US\$5 billion loss per year for wheat, rice, maize and soybean for China, South Korea and Japan. Ghude et al. (2014) estimated India's crop loss for cotton, soybean, rice, and wheat to amount to 6 Mtons with a value of US\$1.3 billion for 2005. These studies, however, may have substantially underestimated the effect O<sub>3</sub> has on crop productivity in Asia due to more sensitivity to O<sub>3</sub> of Asian-grown wheat and rice cultivars (Emberson et al., 2009).

#### Global climate change

The naturally occurring gases of the Earth's atmosphere trap some of the sun's heat in the troposphere. This "greenhouse effect" keeps the Earth's temperature stable. Human activities are producing too much of some of these greenhouse gases (GHGs) (USEPA, 2014e). As a result, the Earth's atmosphere is trapping more of the sun's heat, causing the Earth's average temperature to rise (global warming). Global warming could have significant impacts on human health, agriculture, water resources, precipitation and draughts, forests, wildlife, and coastal areas. Table 4.2 presents some of the properties of GHGs.

4 Controls implemented under the Montreal Protocol have enabled the global community to avoid millions of cases of fatal skin cancer and tens of millions of cases of non-fatal skin cancer and eye cataracts. The U.S. estimates that, by the year 2065, more than 6.3 million skin cancer deaths will have been avoided in that country alone and that efforts to protect the ozone layer will have saved it an estimated US\$4.2 trillion in healthcare costs over the period 1990–2065 (United Nations Development Programme, 2012). Moreover, in 2011, the United States Environmental Protection Agency estimated that more than 22 million Americans born between 1985 and 2100 would avoid suffering from cataracts as a result of the Montreal Protocol (UNEP, 2012).

Table 4.2 Sources and lifetimes of greenhouse gases

Compound	Sources	Lifetime [years]	Reference
CO <sub>2</sub>	Fossil fuel and biomass burning	3–4	Gurjar, et al., 2010
CH <sub>4</sub>	Rice paddies, livestock, natural gas leaks	12	IPCC, 2007
N <sub>2</sub> O	Agriculture, fossil fuel combustion, wastewater management, industrial processes	114	USEPA, 2010; IPCC, 2007
CFCs	Air conditioners, refrigerators, sprays	45–1,700	IPCC, 2007
HFCs		1.4–270	
HCFCs		1.3–17.9	
PFCs		740–50,000	
HFES		0.33–136	
Halons		16–65	

CO<sub>2</sub>: Carbon dioxide; CFCs: Chlorofluorocarbons; HCFCs: Hydrochlorofluorocarbons; HFCs: Hydrofluorocarbons; CH<sub>4</sub>: Methane; PFCs: Perfluorinated compounds; HFES: Fluorinated ethers

### 4.1.3 Link of air pollution and climate change; co-benefits

Many air pollutants such as PM, SO<sub>2</sub>, NO<sub>2</sub>, and CO have the same sources as GHGs. Major GHGs are CO<sub>2</sub>, CH<sub>4</sub>, and O<sub>3</sub>. Ozone formation follows from various chemical transformations such as photochemical oxidation of organic vapors by organic peroxy free radicals in the presence of NO<sub>x</sub>, and CH<sub>4</sub> photo-oxidation via various sequences (Johnston & Kinnison, 1998). Greenhouse gases have to be reduced to mitigate climate change, while air pollutants have to be controlled to avoid direct effects on human and environmental health. Measures to mitigate climate change can reduce air pollution, and actions reducing local and regional air pollution can reduce GHG emissions. There is a need to jointly assess these interrelated policies. The co-control of air pollution and GHG emissions will be cost-effective and create co-benefits (Intergovernmental Panel on Climate Change [IPCC],

2007; Anenberg et al., 2012; Climate and Health Alliance (CAHA), 2012; HEAL, 2012; West et al., 2013; GCEC, 2014). The linkages between air pollution and climate and their relation to health and environmental impacts are depicted in Figure 4.3.

A United Nations Environment Programme/World Meteorological Organization (UNEP/WMO) report has recently compiled scientific studies that have shown that reductions of emissions of short-lived climate pollutants (SLCPs) can potentially prevent millions of premature deaths, prevent the loss of million tons of crops, and reduce global temperatures by 0.5°C [0.2°C–0.7°C] over the next four decades (UNEP/WMO, 2011). Several UNEP-recommended actions to reduce emissions of BC, organic carbon, PM<sub>2.5</sub>, CO, NO<sub>x</sub>, and CH<sub>4</sub> and case studies on co-benefits of linking air pollution and GHG reduction are provided in **Annex III-A and III-B of the Information Sourcebook**.

**Measures to mitigate climate change can reduce air pollution, and actions reducing local and regional air pollution can reduce greenhouse gas emissions. There is a need to jointly assess these interrelated policies.**




Figure 4.3 Linkages between air pollution, climate, and health impacts

Source: Stockholm Environment Institute (SEI), 2008

**Health impact assessment in a clean air action plan is a means of assessing the positive and negative health impacts of existing air pollution abatement policies, strategies, plans and projects.**

### 4.1.4 Health impact assessment in clean air action plans

A clean air action plan (CAAP) is a long-term plan intended to improve air quality and public health by identifying measures to reduce emissions from sectors such as transport, industries, waste deposits, residential burning, among others [see *Guidance Area 5 on Clean air action plans*]. Health impact assessment (HIA) in a CAAP is a means of assessing the positive and negative health impacts of existing air pollution abatement policies, strategies, plans and projects. Many definitions of HIA exist (Birley, 1995; European Centre for Health Policy [ECHP], 1999; Lock, 2000; International Association for Impact Assessment, 2006), the main definition being that of the ECHP: “A combination of procedures, methods and tools by which a policy, program

or project may be judged as to its potential effects on the health of a population, and the distribution of those effects within a population” (WHO, 2007).

Health impact assessment is a valuable tool within a CAAP to develop policies, strategies, programs and projects for cleaner air by providing information for decision makers and addressing policymaking requirements. One of the objectives of an HIA study is to assess whether or not air pollution has an effect on human health. This tool brings policies and people together and involves all stakeholders. It is a proactive process that can mitigate health impacts due to air pollution. The HIA procedure within a CAAP is shown in Figure 4.4. A stepwise process for HIA development and implementation as well as case studies are provided in **Annex III-C of the Information Sourcebook**.




Figure 4.4 The procedure for health impact assessment in clean air action plans

Source: Adapted from WHO, 2015

## 4.2 Stages of health and environmental impact assessment

Strengthening the capacity for health and environmental impact assessment helps shape and define policies for improving air quality. Such capacity also helps in the assessment of the effectiveness of measures to protect human health and the environment. As a starting point, Table 4.3 presents indicators that would aid cities in identifying their current state of AQM development for health and environmental impact assessment.

The following served as key considerations in progressing through the stages:

- Availability of information for estimating health and other impacts
- Processes for estimating health impacts of air pollution
- Capacity for estimating health and other impacts of air pollution
- Presentation of results of health impacts assessment for policy development purposes

Table 4.3 Stages for the assessment of health and other impacts due to air pollution

Stages	Indicators
Underdeveloped	Absence of health surveillance system
	Unavailability of meteorological and air quality databases for emission-exposure-impacts modeling
	Absence of anecdotal observations of and information on health impacts by health authorities
	Lack of capacity for: <ul style="list-style-type: none"> <li>• air pollution monitoring with cheap sensors or sophisticated analyzers with respect to HIA</li> <li>• exposure assessment</li> <li>• health and environmental impact assessment</li> </ul>
Developing	Studies on socio-economic cost of pollution and benefits of pollution control are not available
	Cost effectiveness/cost-benefit analysis is not conducted
	A health surveillance system is being developed
	Meteorological and air quality databases are being developed for emission-exposure-impacts modeling
Developing	Initial observations on health impacts due to air pollution exposure exist
	Capacity is being developed for: <ul style="list-style-type: none"> <li>• air pollution monitoring with cheap sensors or sophisticated analyzers with respect to HIA</li> <li>• exposure assessment</li> <li>• health and environmental impact assessment</li> </ul>
	Studies on socioeconomic cost of pollution and benefits of pollution are not available
	Cost effectiveness/cost-benefit analysis is not conducted

Stages	Indicators
Emerging	<p>A health surveillance system starts to provide reliable data</p> <p>Meteorological and air quality databases are beginning to be established for emission-exposure-impacts modeling</p> <p>Routine observations on health impacts due to air pollution exposure are becoming more and more common</p> <p>Capacity is increasing and regularly enhanced by training for:</p> <ul style="list-style-type: none"> <li>air pollution monitoring with cheap sensors or sophisticated analyzers with respect to HIA</li> <li>exposure assessment</li> <li>health and environmental impact assessment</li> </ul> <p>Limited studies on socioeconomic cost of pollution and benefits of pollution control are available, mostly performed by academic/research institutions</p> <p>Cost-effectiveness/cost-benefit analysis is intermittently conducted by academic/research institutions</p> <p>Presentation of results of HIA, studies on socioeconomic cost of pollution, and cost-effectiveness/cost-benefit analysis considers its use in AQM policy development, implementation, and evaluation</p>
	<p>A health surveillance system is in place and is becoming a basis of HIA due to air pollution</p> <p>Meteorological and air quality databases are routinely used for emission-exposure-impacts modeling</p> <p>Systematic epidemiological studies on health impacts due to air pollution exposure are performed including exposure and health impact assessment studies of major facilities or areas (e.g., schools, hospitals) and of vulnerable populations (e.g., children and the elderly), using rapid assessment techniques</p> <p>Capacity and understanding of issues is increased for:</p> <ul style="list-style-type: none"> <li>air pollution monitoring with cheap sensors or sophisticated analyzers with respect to HIA</li> <li>exposure assessment</li> <li>health and environmental impact assessment</li> </ul> <p>Studies on socioeconomic cost of pollution and benefits of pollution control are becoming available, performed by both academic/research institutions and the government</p> <p>Cost effectiveness/cost benefit analysis is conducted by academic/research institutions and the government</p> <p>Presentation of results of HIA, studies on socioeconomic cost of pollution, and cost-effectiveness/cost-benefit analysis systematically considers its use in AQM policy development, implementation, and evaluation</p>

Stages	Indicators
Fully developed	<p>A health surveillance system makes available reliable data and is always taken as the basis of HIA due to air pollution</p> <p>Meteorological and air quality databases are regulated to be routinely used for emission-exposure-impacts predictions</p> <p>Systematic epidemiological studies on health impacts due to air pollution exposure are performed including exposure and health impact assessment studies of major facilities or areas (e.g., schools, hospitals) and of vulnerable populations (e.g., children and the elderly), using sophisticated assessment techniques</p> <p>Capacity is sustainably enhanced for:</p> <ul style="list-style-type: none"> <li>air pollution monitoring with cheap sensors or sophisticated analyzers with respect to HIA</li> <li>exposure assessment</li> <li>health and environmental impact assessment</li> </ul> <p>Studies on socioeconomic cost of pollution and benefits of pollution control are available, performed by academic/research institutions and the government. There is a process in place for estimating socio-economic costs adapted to local conditions</p> <p>Cost-effectiveness/cost-benefit analyses are routinely performed by academic/research institutions and the government following a localized system for estimating costs and benefits</p> <p>Presentation of results of HIA, studies on socioeconomic cost of pollution, and cost-effectiveness/cost-benefit analysis systematically considers its use in AQM policy development, implementation, and evaluation</p>

### 4.3 Issues and challenges

While a number of Asian cities and countries have developed CAAPs over the last 15 years to address the challenge of air pollution, a number of considerable challenges remain to be resolved, especially in urban areas in Asia. This relates particularly to conducting HIA *vis a vis* air pollution. In addition, the valuation of the benefits of avoiding health impacts within CAAPs is often not performed.

#### Institutional

- **Insufficient human resources and institutional capability within governmental agencies possessing knowledge on the health and environmental impacts of air pollution**, and understanding the linkages between GHG mitigation and air pollution reduction in minimizing health and environmental impacts.
- **Limited understanding of governmental institutions of the importance of studies on health and environmental impacts caused by air pollution.** This understanding needs to get the support of all agencies and other stakeholders in providing data necessary for epidemiological studies.
- **Lack of collaboration, communication and coordination between environmental authorities and health authorities** – preventing agencies from sharing of demographic, monitoring, mortality, and morbidity data needed for epidemiological studies.
- **Lack of understanding among government agencies of the linkage between source and emissions inventory, air quality monitoring, meteorological situation, and health and environmental impacts due to air pollutant exposure.** All these issues are important components of a CAAP, with health and environmental impacts as the most relevant as these could help bring about desired behavior change.
- **Lack of media information and awareness** on: (1) health and environmental impacts of air pollution, and of the health and economic benefits of actions to avoid or minimize the impacts due to air pollution; and (2) impacts of GHG emissions, their linkage to air pollution impacts, and the co-benefits of GHG mitigation and air pollutant emission reduction.

- **Lack of public awareness** that would enable public society groups/non-governmental organizations (NGOs) with interest in combating air pollution and pushing the government to act. These organizations can be interested to launch citizen science projects on air pollution exposure and effect assessment.

#### Management and Technical

- **Limited information on health impacts due to air pollution.** Policy and decision making process are limited and constrained by the poor data and information on health impacts/effects, social and economic effects of the health risks. Limited connection with planning and policy development
- **Only a few Asian countries develop systematic CAAPs** to update or set emission and air quality standards based on health impacts/effects and monitor/enforce the compliance.
- **Limited health impact modeling** and/or short-term and long-term impact estimation by means of epidemiological health impact studies
- **Poor “public health surveillance programs”**—i.e. poor collection of mortality and morbidity information for selected health impacts, lack of cohort and time-series studies.
- **Poor early warning systems to protect the public against air pollution episodes.** These are needed in case of transboundary air pollution (haze episodes), emergency situations due to industrial malfunction, and meteorological events (such as strong inversions).
- **Lack of awareness and studies** on the following:
  - Short-term and long-term health and environmental impacts. Awareness is important to protect the public and the environment against the impacts of air pollution. Local studies are needed to assess the severity of health and environmental impacts.
  - Valuation of health and environmental impacts. Such studies are critical to get information about the benefits of avoiding health and environmental impacts as trade-offs to costs of abatement measures.
  - Linkage between air pollution health impacts and GHG mitigation measures. Awareness of the linkages is critical to understand the interrelationship between air pollution and greenhouse gases.

5 Public health surveillance is defined as the ongoing, systematic collection, analysis and interpretation of health data essential to the planning, implementation, and evaluation of public health practice, closely integrated with the timely dissemination of these data to the appropriate individuals or institutions” (WHO, 1999)

- Risk communication and effective guidance to protect the population from the impacts of heavy pollution
- Valuation of co-benefits of air pollution health impacts and GHG mitigation measures. The co-benefits of simultaneous air pollution reduction and GHG mitigation in terms of health and environmental impacts are usually higher than those of reducing air pollutant emissions and mitigating GHG emissions separately.

#### Financial

In many Asian countries, lack of sufficient governmental funding prevents the development and implementation of health surveillance system and the conduct of epidemiological studies to assess relationships between exposure to air pollutants and health impacts. Funding is especially needed for studies on long-term exposure to air pollutants. Poor communities are constrained financially to introduce and apply protective and adaptation approaches.

In addition, the valuation of the benefits of avoiding health impacts as a consequence of CAAPs is often not performed. The opportunities of tapping into other sources of funding such as public-private-partnerships, grants or loans from international funding agencies, and citizen science or crowdsourcing approaches are often not known or, if known, not explored (Oxford English Dictionary, 2014)<sup>6</sup>.

Despite these challenges, some Asian countries have assessed the local health consequences of air pollution and estimated their economic impact. Health studies in Bangkok and Hong Kong have provided some rationale to take air pollution abatement actions (Lee et al., 2013; Vichit-Vadakan & Vajanpoom, 2011; Hedley, 2009; Vassanadumrongdee & Matsuoka, 2005). The estimated premature deaths, hospital admissions, outpatient visits, and economic loss from air pollution in the Hedley Environmental Index system give such a strong incentive (Box 4.1). The USEPA has collaborated with national experts in several Asian cities to conduct systematic studies on the health and economic impacts of air pollution control policies (e.g. USEPA, 2014h).

#### Box 4.1 Hedley Environmental Index: measuring Hong Kong’s air pollution cost

The Hedley Environmental Index (Hedley Index) closes the gap between generating scientific evidence and communicating the risks in terms of the number of deaths, the number of hospital bed-days, the number of doctor visits, the total economic loss, the loss of tangible cost – hospital admissions, outpatient visits (including travel cost), and work absence, and the loss of healthy life value (Hedley Index, 2014a). Using internationally established exposure-response relationships between air pollutants and health impacts of current levels of pollution, the Hedley Index is acting as a tool to assess the potential public health benefits for strengthened air quality standards in Hong Kong.

On the Hedley Index website (Figure 4.5), data on the five key air pollutant indicators PM<sub>2.5</sub>, PM<sub>10</sub>, NO<sub>2</sub>, SO<sub>2</sub>, and O<sub>3</sub> are reported from three roadside stations and 10 general monitoring stations run by the Hong Kong Environmental Protection Department (EPD). The data is then plotted real-time against the WHO’s short- and long-term Air Quality Guidelines. To delve deeper into the issue, the historical series within a program called “Air Quality Tracker” adds a different function, allowing to review data beginning in 1998, and track the history of each pollutant (Hedley Index, 2014b).

The actual figures in the Hedley Index provide a rather conservative estimate, which does not yet account for air pollution’s effects on vulnerable groups such as pregnant women and young children, the monetary losses of the long-term health burden, or the economic costs of Hong Kong’s lost tourism.

6 Citizen science is also known as crowd science, crowd-sourced science, civic science, volunteer monitoring or networked science.



Figure 4.5 The Hedley Environmental Index Board

According to the Hedley Index, each year, air pollution kills more than 3,000 people in Hong Kong (Kwong, 2012). The Hedley Index is frequently cited and used as a teaching tool by policy researchers, media, and NGOs, and has become a key reference for Hong Kong EPD officials (Ng, 2012; Cheng & Luo, 2009).

The Hedley Index promoted the communication on air quality and its health impact as well as raised the awareness in Hong Kong. On December 30, 2013, Hong Kong EPD launched the Air Quality Health Index to replace the Air Pollution Index, which reports the short-term health risk of air pollution and helps the public take precautionary measures to protect health.

#### 4.4 Roadmap for health and environmental impact assessment

Reliable information on health and environmental impacts is essential for AQM. To implement a roadmap for health and environmental impact assessment within AQM, a first step is to assess where a city is in the AQM developmental stages, then follow the recommendations to progress to the next stage. Table 4.4 gives the recommended steps to implement the roadmap and overcome the challenges outlined above.

Table 4.4 Recommended steps to implement a roadmap for health and other impacts assessment and overcome the challenges of air pollution impacts on health and the environment

Developmental stages	Steps to follow
Underdeveloped	<p><b>Management Process</b></p> <ul style="list-style-type: none"> <li>Start to conduct health and environmental impact assessment (WHO, 2004b; USEPA, 2011b)</li> <li>Start to prepare an air quality monitoring system</li> </ul> <p><b>Technical Process</b></p> <ul style="list-style-type: none"> <li>Learn from international experiences and studies on health impacts and consider the health factors in the policy, plan and strategy development process</li> </ul>
Developing	<p><b>Management Process</b></p> <ul style="list-style-type: none"> <li>Ensure that health risk estimations are used to inform policy makers and are being considered in the policies and plan development</li> <li>Prioritize identification of air pollution control plans and policies that consider impacts on health</li> </ul> <p><b>Technical Process</b></p> <p>Enhance capacity for:</p> <ul style="list-style-type: none"> <li>estimating exposure (Global Atmospheric Pollution Forum (GAPF), 2011)</li> <li>initial estimates of health risks and/or other impacts (WHO, 2004b)</li> <li>an initial health surveillance approach (WHO, 2014d; Nsubuga et al., 2006)</li> </ul>
Emerging	<p><b>Management Process</b></p> <ul style="list-style-type: none"> <li>Use information on health and other impacts to inform policy development</li> <li>Integrate risk assessment into policies</li> </ul> <p><b>Technical Process</b></p> <p>Ensure:</p> <ul style="list-style-type: none"> <li>growing capacity for simplified approaches to estimate exposure (WHO, 2004b)</li> <li>improved understanding of the technical and economic feasibility of major pollution control measures (USEPA, 2011d; 2008b; Reis, 2005)</li> </ul> <p>Enhance capacity for estimates such as:</p> <ul style="list-style-type: none"> <li>the health risks and/or other impacts (WHO, 2006)</li> <li>the economic impacts, based on international studies (Organisation for Economic Co-operation and Development (OECD), 2014)</li> </ul>

Developmental stages	Steps to follow
Maturing	<p><b>Management Process</b></p> <ul style="list-style-type: none"> <li>Establish a robust health statistics database based on a health surveillance system to enable and facilitate health studies</li> <li>Translate study results into policy tools in a transparent manner</li> <li>Develop and implement communication strategies targeted at policy makers and the public</li> <li>Ensure that scenarios for future needs of AQM are developed (USEPA, 2012f)</li> </ul> <p><b>Technical Process</b> Ensure that:</p> <ul style="list-style-type: none"> <li>exposure and HIA studies of major facilities or areas (e.g., schools, hospitals) and of vulnerable populations (e.g., children and the elderly) are conducted (Makri &amp; Stilianakis, 2008)</li> <li>major studies on health impacts are systematically conducted (Rothman &amp; Greenland, 1998)</li> <li>studies on other impacts of air pollution (e.g., on buildings and agriculture) are available (Emberson et al., 2003; Rao et al., 2014)</li> <li>studies on social economic cost of pollution and benefit of pollution control are available (OECD, 2014)</li> <li>cost-effectiveness/cost-benefit analysis are being conducted (USEPA, 2010b; Clean Air for Europe, 2012)</li> <li>emissions inventories and dispersion modeling are further developed (European Environment Agency, 2013)</li> </ul>
Fully developed	<p><b>Management Process</b></p> <ul style="list-style-type: none"> <li>Integrate health risk assessment into policies and strategies</li> <li>Develop and implement communication strategies with policy makers and the public as target audiences</li> </ul> <p><b>Technical Process</b></p> <ul style="list-style-type: none"> <li>Conduct health and environmental impact studies regularly and systematically. Report results to influence policies. Link health and environmental impacts projections based on scenarios with policy changes in emissions-related sectors (e.g., transport). Conduct regularly and systematically exposure and HIA of major facilities or areas (e.g., schools, hospitals) and of vulnerable populations (e.g., children, elderly, and people with existing ailments and/or disabilities). Make available studies on other impacts of air pollution (e.g., on buildings and agriculture) as well as socioeconomic cost of pollution and benefit of pollution control. Conduct cost-effectiveness/cost-benefit analysis.</li> <li>Develop regulatory air quality simulation models to support the policymaking process</li> </ul>

Municipal, provincial and national governments should recognize the need to consider health and environmental impacts as an important ingredient of AQM. In fact, if there were no impacts of air pollution, there would also be no need for action. In developing Asian countries, it is important to strengthen the political will and understanding of the social and economic costs of air pollution, which often surpass the costs of control measures [See *Guidance Area 6 on Governance*]. A strong governmental response is needed to mitigate the health and environmental effects of air pollution (Box 4.2). It is also important to strengthen the linkages among source inventory, emission, air quality monitoring, meteorological situation, and health and environmental impacts due to air pollutant exposure. To better understand these linkages, it is necessary to involve all relevant stakeholders in the development of action plans for health and environmental impact assessment as an integral part of CAAPs [See *Guidance Area 5 on Clean air action plans*]. Similarly, the collaboration, communication, and coordination between environmental authorities and health authorities on the municipal, provincial and national levels should be strengthened. Enhancing the institutional capability with respect to AQM issues through sustainable education and training and provision of sufficient human resources is also significant [See *Guidance Area 6 on Governance*].

Sufficient financial resources should be allocated for the following health and environmental impact studies. If reliable emissions and meteorological data exist, models can be used to estimate exposure [See *Guidance Area 2 on Emissions inventories and modeling*] and – with the application of established exposure-response relationships – corresponding health environmental impacts. This would be the low-cost option because appropriate models can be downloaded free of charge e.g. the USEPA website. If a health surveillance system is in place, epidemiological studies can be performed but the cost of such studies can be high to very high depending on the type of study, the number of cases and the planned duration of a study.

Financial resources should be made available for:

- emission-exposure-impact studies
- a health surveillance system (if it does not exist)
- short-term and long-term health and environmental studies
- studies on the valuation of health and environmental impacts
- studies on the linkage between air pollution health impacts and GHG mitigation measures
- studies on the valuation of co-benefits of air pollution health impacts and GHG mitigation measures

**Municipal, provincial and national governments should recognize the need to consider health and environmental impacts as an important ingredient of air quality management. In fact, if there were no impacts of air pollution, there would also be no need for action. In developing Asian countries, it is important to strengthen the political will and understanding of the social and economic costs of air pollution, which often surpass the costs of control measures.**

## Box 4.2 Responding to the health effects of air pollution in India

Outdoor air concentrations of various air pollutants in Indian cities are continuing to be a major health concern in India because of their persistent high levels. The impact of long-term exposure to urban air pollution on the respiratory and other organ systems of the body was studied in Kolkata and Delhi, two highly polluted megacities in the country (Ray & Lahiri, 2010). Moreover, the health impacts of vehicular pollution during 2007 to 2010 have been investigated (Ray & Lahiri, 2012). Compared to rural controls, the urban population had a significantly higher prevalence of upper and lower respiratory symptoms, bronchial asthma, and lung function deficits. The adverse health consequences in the urban population were positively associated with PM<sub>10</sub> levels in ambient air and with personal exposures to benzene. Long-term exposures to high levels of urban air pollution in these cities are adversely affecting the physical and mental health of citizens, especially of children and the elderly. In Kolkata, long-term exposure to air pollution arises mostly from vehicular exhausts.

Another study estimated the number of premature deaths in 14 Indian cities (Nema & Goyal, 2010). Based on the PM<sub>10</sub> concentration levels of 2001, among the metro cities, the highest number of mortality cases is observed in Delhi (4889), followed by Kolkata (4303), Mumbai (1959), and Chennai (1272). India has set up institutions to respond to the air pollution and its health impacts. The Ministry of Environment, Forest, and Climate Change (MoEF) is the nodal agency in the Central Government for overseeing the implementation of India's environmental policies and programs. The MoEF initiated environmental epidemiological studies in different areas of indoor and outdoor pollution to identify and develop programs that shall create a database and suggest environmental mitigation measures.

The Central Pollution Control Board (CPCB) under MoEF is responsible for planning and executing comprehensive nationwide programs for the prevention and control of air pollution. The CPCB is executing a nationwide National Air Quality Monitoring Program (NAMP). The NAMP is covering 545 operating stations spread over covering 225 cities/towns in 26 States and five Union Territories. A system for collecting real-time online data has been established and publishes air quality data from 35 continuously monitoring stations operated by various agencies (MoEF, 2014). The CPCB has stipulated National Ambient Air Quality Standards (NAAQS) since 1982. The NAAQS are based on health criteria and follow a land use based approach. The NAAQS have been revised in November 2009 in consultation with the civil society and experts, for 12 pollutants including SO<sub>2</sub>, NO<sub>2</sub>, PM<sub>10</sub>, PM<sub>2.5</sub>, O<sub>3</sub>, Pb, As, Ni, CO, NH<sub>3</sub>, benzene, and B(a)P. These standards/limits provide a legal framework for the control of air pollution and the protection of public health.

In January 2014, a Steering Committee on health related issues on Air Pollution was formed, with a view towards framing an action plan for mitigating the adverse health impacts of indoor and outdoor air pollution.

In June 2015 the Government of India launched the air quality index in New Delhi and urged the people to change their lifestyle in order to help protect the environment (Hindustan Times, 2015).

As: Arsenic; B(a)P: Benzo(a)pyrene; NH<sub>3</sub>: Ammonia; Ni: Nickel; Pb: Lead

Source: CPCB, 1998; MOEF, 2014

## 4.5 References

- Anenberg, S., Schwartz, J., Shindell, D., Amann, A., Faluvegi, G., Klimont, Z., Janssens-Maenhout, G., Pozzoli, L., Van Dingenen, R., Vignati, E., Emberson, L., Muller, N.Z., West, J.J., Williams, M., Demkine, V., Hicks, W.K., Kuylenstierna, J., Raes, F., Ramanathan, V., 2012. Global air quality and health co-benefits of mitigating near-term climate change through methane and black carbon emission controls. *Environmental Health Perspectives*, 120: 831-839.
- Avnery, S., Mauzerall, D.L., Liu, J., & Horowitz, L.W. (2011). Global crop yield reductions due to surface ozone exposure: Year 2000 crop production losses and economic damage. *Atmospheric Environment*, 45: 2284–2296. Retrieved from <https://www.princeton.edu/~mauzeral/papers/Avnery%20Mauzerall%20etal%20ag%202000%20AE%202011%20Global%20Crop%20Yield.pdf>.
- Birley, M. (1995). *The health impact assessment of development projects*. London, U.K.: Her Majesty's Stationery Office.
- Brauer, M., Amann, M., Burnett, R.T., Cohen, A., Dentener, F., Ezzati, M., Henderson, S.B., Krzyzanowski, M., Martin, R.V., Van Dingenen, R., Van Donkelaar, & A., Thurston, G.D. (2012). Exposure assessment for estimation of the global burden of disease attributable to outdoor air pollution. *Environmental Science and Technology*, 46: 652–660. doi:10.1021/es2025752
- Brook, R.D., Rajagopalan, S., Pope III, C.A., Brook, JR, Mittleman, M.A., Peters, A., Siscovick, D., Smith, S.C., Whitsel, L., & Kaufman, J.D. (2010). *Particulate matter air pollution and cardiovascular disease: An update to the scientific statement from the American Heart Association* (Statement from the American Heart Association). *Circulation* 121: 2331–2378. doi: 10.1021/es2025752
- Central Pollution Control Board, India. (1998). *Ambient air quality standard for ammonia*. New Delhi, India: Author. Retrieved from <http://envfor.nic.in/legis/air/gsr935%28e%29.htm>.
- Cheng, E. & Luo, R. (2009). *Hedley's Environmental Index sheds new light on Hong Kong's air pollution*. Washington, D.C.: Wilson Center. Retrieved from <http://www.wilsoncenter.org/publication/hedley-environmental-index-sheds-new-light-hong-kongs-air-pollution>
- Chuang, K.J., Chan, C.C., Su, T.C., Lee, C.T., & Tang, C.S. (2007). The effect of urban air pollution on inflammation, oxidative stress, coagulation, and autonomic dysfunction in young adults. *American Journal of Respiratory and Critical Care Medicine*, 176:370–376. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/17463411>
- Clean Air for Europe. (2012). *Clean Air for Europe – Cost benefit analysis*. Retrieved from <http://www.cafe-cba.org/>.
- Climate and Health Alliance and the Climate Institute. (2012). *Our uncashed dividend. The health benefits of climate action*. Retrieved from [http://caha.org.au/wp-content/uploads/2012/08/OurUncashedDividend\\_CAHAandTCI\\_August2012.pdf](http://caha.org.au/wp-content/uploads/2012/08/OurUncashedDividend_CAHAandTCI_August2012.pdf)
- Cyclopedic Medical Dictionary. (1997). *Taber's cyclopedic medical dictionary*. (Edition 18). F.A. Philadelphia, U.S.A., Davis Company, Philadelphia.
- Emberson, L., Ashmore, M., Murray, & F., (Eds.). (2003). *Air pollution impacts on crops and forests. A global assessment*. London, U.K.: Imperial College Press.
- Emberson, L.D., Büker, P., Ashmore, M.R., et al. (2009). A comparison of North American and Asian exposure-response data for ozone effects on crop yields. *Atmospheric Environment*, 43: 1945–1953.



European Centre for Health Policy. (1999). *Health impact assessment: main concepts and suggested approach* (Gothenburg consensus paper). Copenhagen, Denmark: WHO Regional Office for Europe.

European Environmental Agency. (2013). *EMEP/EEA air pollutant emission inventory guidebook*. Retrieved from <http://www.eea.europa.eu/themes/air/emep-eea-air-pollutant-emission-inventory-guidebook>

Global Atmospheric Pollution Forum & Schwela, D. (2011). *GAP Forum air pollution monitoring manual*, York, U.K.: Stockholm Environment Institute. Retrieved from <http://sei-international.org/rapid/gapforum/html/projects.php>

Global Atmospheric Pollution Forum. (2012). *The Forum air pollutant emissions inventory preparation manual* (Version 5). Retrieved from <http://www.sei-international.org/gapforum/tools.php>

Ghude, S.D., Jena, C., Chate, D.M., Beig, G., Pfister, G.G., Kumar, R., & Ramanathan, V. (2014). Reductions in India's crop yield due to ozone. *Geophysical Research Letters*, 41: 5685–5691. Retrieved from <http://www.ramanathan.ucsd.edu/files/pr204.pdf>

Global Commission on the Economy and Climate Website. (2014). The new climate economy. Chapter 4. Energy. Better energy, better climate. *The Global Commission on the Economy and Climate*. Retrieved from: <http://newclimateeconomy.report/energy/>.

Gurjar B.R., L. T., Molina, C.S. P., & Ojha. (2010). *Air Pollution: Health and Environmental Impacts*. Boca Raton: CRC press.

Health and Environmental Alliance. (2012). *The health benefits of greenhouse gas reduction policies*. Retrieved from [http://www.google.de/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&ved=0CClQFjAAahUKewi99Y7pvpbGAhXGiywKHdjvAL0&url=http%3A%2F%2Fwww.env-health.org%2FIMG%2Fpdf%2Fheal\\_background\\_paper\\_climate\\_co-benefits\\_en.pdf&ei=BEiBVb3jEsaXsgHY34PoCw&usq=AFQjCNFIFNnd\\_vl6JL7p6JGsX539ISBHUA&bvm=bv.96041959,d.bGg&cad=rja](http://www.google.de/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&ved=0CClQFjAAahUKewi99Y7pvpbGAhXGiywKHdjvAL0&url=http%3A%2F%2Fwww.env-health.org%2FIMG%2Fpdf%2Fheal_background_paper_climate_co-benefits_en.pdf&ei=BEiBVb3jEsaXsgHY34PoCw&usq=AFQjCNFIFNnd_vl6JL7p6JGsX539ISBHUA&bvm=bv.96041959,d.bGg&cad=rja)

Health Effects Institute. (2013). *National Particle Component Toxicity Initiative report on cardiovascular effects* (Research Report 178). Retrieved from <http://pubs.healtheffects.org/getfile.php?u=946>.

Hedley, A.J., 2009. *Air pollution and public health - The current avoidable burden of health problems, community costs and harm to future generations*. CB (1) 733/08-09(02). Hong Kong: University of Hong Kong.

Hedley Environmental Index. (2014a). *Cumulative loss due to air pollution*. Retrieved from <http://hedleyindex.sph.hku.hk/html/en/>

Hedley Environmental Index. (2014b). Hedley Environmental Index. *Cumulative loss due to air pollution*. Retrieved from <http://hedleyindex.sph.hku.hk/html/en/chart.php>

Hirota, N. & Martin, J.G. (2013). Mechanisms of airway remodeling. *Chest* 144: 1026–1032. Retrieved from [http://publications.chestnet.org/data/Journals/CHEST/927499/chest\\_144\\_3\\_1026.pdf](http://publications.chestnet.org/data/Journals/CHEST/927499/chest_144_3_1026.pdf).

Hogg, J.C., Chu, F., Utokaparch, S., Woods, R., Elliott, W.R., Buzatu, L., Cherniack, R.M., Rogers, R.M., Scuirba, F.C., Harvey, O., & Pare, P.D., (2004). The nature of small-airway obstruction in chronic obstructive pulmonary disease. *New England Journal of Medicine*, 350(26): 2645–2653. doi: 10.1056/NEJMoa032158

Modi launches air quality index, wants change in lifestyle to save environment. (2015, June 4). *Hindustan Times*. Retrieved from <http://www.hindustantimes.com/india-news/pm-modi-launches-air-quality-index-says-environment-and-development-can-go-hand-in-hand/article1-1334293.aspx>

International Association for Impact Assessment. (2006). *Health impact assessment –international best practices*. Retrieved from: <http://www.iaia.org/publicdocuments/special-publications/SP5.pdf?AspxAutoDetectCookieSupport=1>.

International Agency for Research on Cancer. (1987). *Summaries & evaluations: Benzene (Group 1)* (IARC Monographs on the Carcinogenicity of Chemicals to Humans, Supplement 7). Retrieved from <http://www.inchem.org/documents/iarc/suppl7/benzene.html>

International Agency for Research on Cancer. (2009a). A review of human carcinogens–Part F: Chemical agents and related occupations (Special Report: Policy). *Lancet Oncology*, 10: 1143–1144.

International Agency for Research on Cancer. (2009b). A review of human carcinogens –Part C: metals, arsenic, dusts and fibers (Special Report: Policy). *Lancet Oncology*, 10: 453-454.

International Agency for Research on Cancer (2012a). Chemical agents and related occupations: A review of human carcinogens. *IARC Monographs on the evaluation of carcinogenic risks to humans*, Volume 100F. Nancy, France: International Agency for Research on Cancer. Retrieved from <http://monographs.iarc.fr/ENG/Monographs/vol100F/mono100F.pdf>

International Agency for Research on Cancer. (2012b). Arsenic, metals, fibres, and dusts. A review of human carcinogens. IARC Monographs on the evaluation of carcinogenic risks to humans, Volume 100 C. Nancy, France: International Agency for Research on Cancer. Retrieved from <http://monographs.iarc.fr/ENG/Monographs/vol100C/index.php>

Intergovernmental Panel on Climate Change. (2007). Changes in atmospheric constituents and in radiative forcing (Contribution of working group I to the Fourth Assessment Report of the IPCC). In Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B., Tignor, & M., Miller, H.L. (Eds.). *The Physical Science Basis*. Retrieved from [http://www.ipcc.ch/publications\\_and\\_data/ar4/wg1/en/contents.html](http://www.ipcc.ch/publications_and_data/ar4/wg1/en/contents.html)

Intergovernmental Panel on Climate Change. (2007). *Climate change 2007* (Contribution of working group III to the Fourth Assessment Report of the IPCC). Retrieved from [https://www.ipcc.ch/publications\\_and\\_data/ar4/wg3/en/tssts-ts-7-6-co-benefits-of.html](https://www.ipcc.ch/publications_and_data/ar4/wg3/en/tssts-ts-7-6-co-benefits-of.html)

Institute of Occupational Medicine and Searl, A., (2004). *A review of the acute and long-term impacts of exposure to nitrogen dioxide in the United Kingdom* (Research Report TM/04/03). Edinburgh, UK: IOM. Retrieved from: [http://www.iom-world.org/pubs/iom\\_tm0403.pdf](http://www.iom-world.org/pubs/iom_tm0403.pdf)

Janssen, N.A.H., Gerlofs-Nijland, M.E., Lanke, T., Salonen, R.O., Cassee, F., Hoek, G., Fischer, P., Brunekreef, B., & Krzyzanowski, M. (2012). *Health effects of black carbon*. Copenhagen, Denmark: WHO Regional Office for Europe. Retrieved from [http://www.euro.who.int/\\_\\_data/assets/pdf\\_file/0004/162535/e96541.pdf](http://www.euro.who.int/__data/assets/pdf_file/0004/162535/e96541.pdf)

Johnston, H., & Kinnison, D. (1998). Methane photooxidation in the atmosphere: Contrast between two methods of analysis. *Journal of Geophysical Research*, 103 (D17): 21967–21984. doi: 10.1029/98JD01213.

Kwong, S.Y., (2012). *New air quality app provides 24-hour forecast and clearer picture of pollution health risks*. Retrieved from <http://www.hongkongcan.org/eng/2013/10/new-air-quality-app-provides-24-hour-forecast-and-clearer-picture-of-pollution-health-risks/>

Lanata, C.F., Rudan, I., Boschi-Pinto, C., Tomaskovic, L., Cherian, T., Weber, M., & Campbell, H. (2004). Methodology and quality issues in epidemiological studies of acute lower respiratory infections in children in developing countries. *International Journal of Epidemiology*, 33(6): 1362–1372. doi: 10.1093/ije/dyh229.

Lee, S.L., Tinsley, H., Chau, J., Lai, H.K., Thach, T.O., Ip, P., Chow, C.B., & Hedley, A.J., 2013. Review of outdoor air pollution and child health in Hong Kong. *Hong Kong Journal of Pediatrics* (new series) 18: 69-76.

Lim, S.S., Vos, T., Flaxman, A.D., Danaei, G., Shibuya, K., Adair-Rohani, H., Ezzati, M. (2012). A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*, 380(9859): 2224–2260.

Long, C.M., Nascarella, M.A., & Valberg, P.A. (2013). Carbon black vs. black carbon and other airborne materials containing elemental carbon: Physical and chemical distinctions. *Environmental Pollution*, 181: 271–286.

Loomis, D., Grosse, Y., Lauby-Secretan, B., El Ghissassi, F., Bouvard, V., Benbrahim-Tallaa, L., Guha, N., Baan, R., Mattock, H., & Straif, K. (2013). The carcinogenicity of outdoor air pollution (On behalf of the International Agency for Research on Cancer Monograph Working Group IARC). *Lancet Oncology* 14(13): 1262–1263. doi: 10.1016/S1470-2045(13)70487-X.

Makri, A., Stilianakis, N.I. (2008). Vulnerability to air pollution effects. *International Journal of Hygiene and Environmental Health*, 211(3-4): 326–336.

Martin, R. (2008). Ambient ammonia – air quality and health impacts. Retrieved from [https://www.google.de/search?q=IARC+nickel+unit+risk&ie=utf-8&oe=utf-8&gws\\_rd=cr&ei=5TWWVbSbJpXUNf\\_gaAG#q=US+EPA+ammonia+standard](https://www.google.de/search?q=IARC+nickel+unit+risk&ie=utf-8&oe=utf-8&gws_rd=cr&ei=5TWWVbSbJpXUNf_gaAG#q=US+EPA+ammonia+standard)

Ministry of Environment, Forest and Climate Change, India. (2014). *Annual report of 2013-14 of Government of India*. Retrieved from [http://www.moef.nic.in/sites/default/files/annual\\_report/AR-2013-14-Eng.pdf](http://www.moef.nic.in/sites/default/files/annual_report/AR-2013-14-Eng.pdf)

Nema, P., & Goyal, S.K., (2010). Estimation of Health Impacts due to PM<sub>10</sub> in Major Indian Cities. In Gurjar, B.R., Molina, L.T. Ojha, & C.S.P., (Eds.), *Air Pollution: Health and Environmental Impacts*. Boca Raton, U.S.A.: CRC Press.

Ng, S.K.W., (2012). *Hedley Environmental Index, May 7, 2012*. Retrieved from [http://cleanairinitiative.org/portal/sites/default/files/presentations/03\\_-\\_Simon\\_Ng\\_-\\_Hedley\\_Index\\_Da\\_Li\\_Zhi\\_Shu\\_.pd](http://cleanairinitiative.org/portal/sites/default/files/presentations/03_-_Simon_Ng_-_Hedley_Index_Da_Li_Zhi_Shu_.pd), [http://climpol.iass-potsdam.de/sites/climpol/files/wysiwyg/files/ig-wg\\_slcpc.pdf](http://climpol.iass-potsdam.de/sites/climpol/files/wysiwyg/files/ig-wg_slcpc.pdf)

Nsubuga, P., White, M.E., Thacker, S.B., Anderson, M.A., Blount, S.B., Broome, C.V., Chiller, T.M., Imtiaz, V.E.R., Sosin, D., Stroup, D.F., Tauxe, R.V., Vijayaraghavan, M., & Trostle, M. (2006). Public health surveillance: A tool for targeting and monitoring. In Jameson, D.T., Breman, J.G., Measham, A.R., Alleyne, G., Claeson, M., Evans, D.B., Jha, P., Mills, A., Musgrove, P., (Eds.), *Disease control priorities in developing countries* (2nd edition). Washington, D.C.: The World Bank.

Organisation for Economic Co-operation and Development. (2014). *The costs of air pollution: Health impacts of road transport*. Retrieved from [http://www.keepeek.com/Digital-Asset-Management/oced/environment/the-cost-of-air-pollution\\_9789264210448-en#page1](http://www.keepeek.com/Digital-Asset-Management/oced/environment/the-cost-of-air-pollution_9789264210448-en#page1)

Oxford English Dictionary. (2014). New word notes. Retrieved from <http://public.oed.com/the-oed-today/recent-updates-to-the-oed/june-2014-update/new-words-notes-june-2014/>

Paulot, F., Jacob, D.J., (2014). Hidden costs of U.S. agricultural export: Particulate matter from ammonia emissions. *Environmental Science and Technology*, 48(2): 903-908. Abstract retrieved from <http://pubs.acs.org.ezproxy.york.ac.uk/doi/abs/10.1021/es4034793>

Phinney, L., Beauchamp, S., Waugh, D., & Tordon, R. (2004). *Environmental impacts of air pollution*. Retrieved from <http://www.novascotia.ca/nse/air/docs/phinney-environmentalimpacts.pdf>.

Rai, R., & Agrawal, M., (2014). Assessment of competitive ability of two Indian wheat cultivars under ambient O<sub>3</sub> at different developmental stages. *Environmental Science and Pollution Research International*, 21(2):1039–53. doi: 10.1007/s11356-013-1981-6.

Ramanathan V., Crutzen P.J., Kiehl J.T., & Rosenfeld D. (2001). Aerosols, climate, and the hydrological cycle. *Science*, 294: 2119–2124, doi:10.1126/science.1064034.

Ramanathan V. & Ramana, M.V. (2003). Atmospheric brown clouds. *Environmental Management*. December 2003, 28–33.

Ramanathan, V., Agrawal, M., Akimoto, H., Aufhammer, M., Devotta, S., Emberson, L., Hasnain, S.I., Iyengararasan, M., Jayaraman, A., Lawrence, M., Nakajima, T., Oki, T., Rodhe, H., Ruchirawat, M., Tan, S.K., Vincent, J., Wang, J.Y., Yang, D., Zhang, Y.H., Autrup, H., Barregard, L., P. Bonasoni, Brauer, M., Brunekreef, B., Carmichael, G., Chung, C.E., Dahe, J., Feng, Y., Fuzzi, S., Gordon, T., Gosain, A.K., Htun, N., Kim, J., Mourato, S., Naeher, L., Navasumrit, P., Ostro, B., Panwar, T., Rahman, M.R., Ramana, M.V., Rupakheti, M., Settachan, D., Singh, A.K., Helen, G.S., Tan, P.V., Viet, P.H., Yinlong, J., Yoon, S.C., Chang, W.-C., Wang, X., Zelikoff, & J., Zhu, A. (2008). *Atmospheric brown clouds: Regional assessment report with focus on Asia*. Nairobi, Kenya: United Nations Environment Programme. Retrieved from <http://www.rrcap.ait.asia/abc/impact/index.cfm>

Rao, N.V., Rajasekhar, M., & Rao, G.C. (2014). Detrimental effect of air pollution, corrosion on building materials and historical structures. *American Journal of Engineering Research*, 3(3): 359-364. Retrieved from [www.ajer.org/papers/v3\(3\)/ZT33359364.pdf](http://www.ajer.org/papers/v3(3)/ZT33359364.pdf)

Ray, M.R., & Lahiri, T. (2010). Health Effects of Urban Air Pollution in India. In Gurjar, B.R., Molina, L.T., Ojha, C.S. P. (Eds.), *Air Pollution: Health and Environmental Impacts*. Boca Raton, U.S.A.: CRC Press.

Ray, M.R., Lahiri, T., (2012). *Air pollution and its effects on health – Case studies, India*. Retrieved from [http://www.theicct.org/sites/default/files/MKRay\\_0.pdf](http://www.theicct.org/sites/default/files/MKRay_0.pdf)

Reis, S. (2005). Costs of air pollution control. Analyses of emission control options for ozone abatement strategies. New York, U.S.A.: Springer Press.

Risom, L., Møller, P., Loft, S. (2005) Oxidative stress-induced DNA damage by particulate air pollution. *Mutation Research*, 592(1-2): 119–137. doi: 10.1016/j.mrfmmm.2005.06.012.

Riva, D. R., Magalhaes, C.B., Lopes, A.A., Lançiv, T., Mauad, T., Malm, O., Valen V, S.S., Saldiva, P.H., Faffe, D.S., & Zin, W.A. (2011). Low dose of fine particulate matter (PM<sub>2.5</sub>) can induce acute oxidative stress, inflammation and pulmonary impairment in healthy mice. *Inhalation Toxicology*, 23: 257–267. doi: 10.3109/08958378.2011.566290.

Roth, D.E., Caulfield, L.E., Ezzati, M., & Black, R.E. (2008). Acute lower respiratory infections in childhood: Opportunities for reducing the global burden through nutritional interventions. *Bulletin of the World Health Organization*, 86; 356-364. Retrieved from: <http://www.who.int/bulletin/volumes/86/5/07-049114.pdf?ua=1>

Rothman, K.J., & Greenland, S. (1998). *Modern epidemiology* (Second edition). Philadelphia: Lippincott Williams & Wilkins.

United Kingdom website. (2014). Impacts of air pollution & acid rain on wildlife. Retrieved from: <http://www.air-quality.org.uk/17.php>.

United Kingdom website. (2014a). *What is acid rain?* Retrieved from: <http://www.air-quality.org.uk/01.php>

---

United Kingdom website. (2014b). *Impacts of Air Pollution & Acid Rain on Wildlife*. Retrieved from: <http://www.air-quality.org.uk/17.php>.

United Nations Development Programme. (2012). *Information note on 25th anniversary of the Montreal Protocol – UNDP activities under the protocol*. Retrieved from: [http://www.undp.org/content/undp/en/home/ourwork/environmentandenergy/focus\\_areas/ozone\\_and\\_climate/UNDP25thMP.html](http://www.undp.org/content/undp/en/home/ourwork/environmentandenergy/focus_areas/ozone_and_climate/UNDP25thMP.html)

United Nations Environment Programme. (2010). *The 2010 assessment of the Scientific Assessment Panel*. Ozone Secretariat. Retrieved from [http://ozone.unep.org/new\\_site/en/scientific\\_assessment\\_2010.php](http://ozone.unep.org/new_site/en/scientific_assessment_2010.php)

United Nations Environment Programme. (2011). *The Montreal Protocol on Substances that Delete the Ozone Layer*. Retrieved from [http://ozone.unep.org/new\\_site/en/montreal\\_protocol.php](http://ozone.unep.org/new_site/en/montreal_protocol.php).

United Nations Environment Programme. (2012). *Key achievements of the Montreal Protocol to date*. Retrieved from [http://www.unep.org/ozonaction/Portals/105/documents/Ozone\\_Day\\_2012/Key%20achievements%20of%20the%20Montreal%20Protocol%202012%20low%20res.pdf](http://www.unep.org/ozonaction/Portals/105/documents/Ozone_Day_2012/Key%20achievements%20of%20the%20Montreal%20Protocol%202012%20low%20res.pdf)

United States Environmental Protection Agency. (2007). *Lead: Human exposure and health risk assessments for selected case studies. Volume I. Human exposure and health risk assessments –full scale*. EPA-452/R-07-014a. Retrieved from [http://www.epa.gov/ttn/naaqs/standards/pb/s\\_pb\\_2010\\_rea.html](http://www.epa.gov/ttn/naaqs/standards/pb/s_pb_2010_rea.html).

United States Environmental Protection Agency. (2008a). *Integrated science assessment for oxides of nitrogen and sulfur – Ecological criteria*. Retrieved from <http://cfpub.epa.gov/ncea/cfm/recorddisplay.cfm?deid=201485>

United States Environmental Protection Agency. (2008b). *Cost analysis models/tools*. Retrieved from <http://www.epa.gov/ttn/ecas/costmodels.html>.

United States Environmental Protection Agency. (2009). *Integrated science assessment for particulate matter*. EPA/600/R-08/139F. Retrieved from <http://www.epa.gov/blackcarbon/effects.html>.

United States Environmental Protection Agency. (2010a). *Methane and Nitrous Oxide Emissions from Natural Sources*. Retrieved from <http://epa.gov/climatechange/ghgemissions/gases/n2o.html>

United States Environmental Protection Agency. (2010b). *The benefits and costs of the Clean Air Act: 1990 to 2020* (Revised draft report, August 2010). Retrieved from <http://www.epa.gov/cleanairactbenefits/aug10/fullreport.pdf>

United States Environmental Protection Agency. (2011a). *Basic information on stratospheric ozone depletion*. Retrieved from: <http://www.epa.gov/ozone/science/basicinfo.html>.

United States Environmental Protection Agency. (2011b). *Health and environmental assessment –resources: Organizations and agencies*. Retrieved from [http://www.epa.gov/oaqps001/aqportal/management/links/assessment\\_resources\\_org.htm](http://www.epa.gov/oaqps001/aqportal/management/links/assessment_resources_org.htm)

United States Environmental Protection Agency. (2011c). *Air quality management portal*. Retrieved from <http://www.epa.gov/airquality/aqportal/>

United States Environmental Protection Agency. (2011d). *Control strategies*. Retrieved from [http://www.epa.gov/airquality/aqportal/management/control\\_strategies.htm](http://www.epa.gov/airquality/aqportal/management/control_strategies.htm)

---

United States Environmental Protection Agency. (2011e). *Cancer incidence: Report on the environment*. Retrieved from <http://cfpub.epa.gov/eroe/index.cfm?fuseaction=detail.viewMeta&ch=49&IShowInd=0&subtop=209&lv=list.listByChapter&r=219651>.

United States Environmental Protection Agency. (2012a). *Black Carbon effects on public health and the environment. In Report to Congress on Black Carbon*, pp. 67-84. Retrieved from [www.epa.gov/blackcarbon/2012report/Chapter3.pdf](http://www.epa.gov/blackcarbon/2012report/Chapter3.pdf). Also see: [www.epa.gov/blackcarbon/](http://www.epa.gov/blackcarbon/)

United States Environmental Protection Agency. (2012b). *Effects of acid rain*. Retrieved from <http://www.epa.gov/acidrain/effects/index.html>

United States Environmental Protection Agency. (2012c). *Visibility*. Retrieved from <http://www.epa.gov/oar/visibility/>

United States Environmental Protection Agency. (2012d). *Effects of acid rain – Forests*. Retrieved from <http://www.epa.gov/acidrain/effects/forests.html>.

United States Environmental Protection Agency. (2012e). *Ecosystem effects*. Retrieved from <http://www.epa.gov/groundlevelozone/ecosystem.html>

United States Environmental Protection Agency. (2012f). *Scenarios for the future of air quality: Planning and analysis in an uncertain world*. Retrieved from [http://www.epa.gov/ttnecas1/workingpapers/WorkingPaperFutureScenarios\\_07Nov12.pdf](http://www.epa.gov/ttnecas1/workingpapers/WorkingPaperFutureScenarios_07Nov12.pdf)

United States Environmental Protection Agency. (2013a). *Integrated science assessment for ozone and related photochemical oxidants*. Retrieved from <http://www.epa.gov/ncea/isa/>

United States Environmental Protection Agency. (2013b). *Integrated science assessment for sulfur oxides – health criteria* (Final report dated September 2008). Retrieved from <http://cfpub.epa.gov/ncea/isa/recorddisplay.cfm?deid=198843#Download>

United States Environmental Protection Agency. (2013c). *Arsenic compounds – hazard summary* (Created in 1992; revised in December 2012). Retrieved from <http://www.epa.gov/ttnatw01/hlthef/arsenic.html>

United States Environmental Protection Agency. (2013d). *Nickel compounds – hazard summary* (Created in 1992; revised in January 2000). Retrieved from <http://www.epa.gov/ttnatw01/hlthef/nickel.html>

United States Environmental Protection Agency. (2014a). *Health effects in the general population*. Retrieved from <http://www.epa.gov/apti/ozonehealth/population.html>.

United States Environmental Protection Agency. (2014b). *Nitrogen dioxide–health*. Retrieved from <http://www.epa.gov/oaqps001/nitrogenoxides/health.html>.

United States Environmental Protection Agency. (2014c). *Integrated science assessment for oxides of nitrogen – health criteria* (Final report). Retrieved from <http://cfpub.epa.gov/ncea/cfm/recorddisplay.cfm?deid=194645#Download>

United States Environmental Protection Agency. (2014d). *Sulfur dioxide–health*. Retrieved from <http://www.epa.gov/airquality/sulfurdioxide/health.html>

---

United States Environmental Protection Agency. (2014e). *Overview of greenhouse gases*. Retrieved from <http://www.epa.gov/climatechange/ghgemissions/gases.html>

United States Environmental Protection Agency. (2014f). *Lead in air: Health*. Retrieved from <http://www.epa.gov/airquality/lead/health.html>

United States Environmental Protection Agency. (2014g). *National ambient air quality standards (NAAQS)*. Retrieved from <http://www.epa.gov/air/criteria.html>

United States Environmental Protection Agency. (2014h). *EPA Collaboration with India. Improving air quality*. Retrieved from <http://www2.epa.gov/international-cooperation/epa-collaboration-india>

United States Geological Survey. (2014). *Eutrophication*. Retrieved from <http://toxics.usgs.gov/definitions/eutrophication.html>

Vassanadumrongdee, S., & Matsuoka, S. (2005). Risk perception and value of a statistical life for air pollution and traffic accidents: Evidence from Bangkok, Thailand. *Journal of Risk and Uncertainty*, 30(3): 261-287.

Vichit-Vadakan, N., & Vajanpoom, N. (2011). Health impact from air pollution in Thailand: Current and future challenges. *Environmental Health Perspectives*, 119(5): A197-A198.

Wang, X., & Mauzerall, D. L. (2004). Characterizing distributions of surface ozone and its impact on grain production in China, Japan and South Korea: 1990 and 2020. *Atmospheric Environment*, 38: 4383-4402.

West, J.J., Smith, S.J., Silva, R.A., Nak, V., Zhang, Y., Adelman, Z., Fry, M.M., Anenberg, S., & Horowitz, L.W. (2013). Co-benefits of mitigating greenhouse gas emissions for future air quality and human health. *Nature Climate Change*, 3: 885-889. doi:10.1038/nclimate2009.

World Health Organization. (1993). *Benzene Environmental Health Criteria 150*. Geneva, Switzerland: International Programme on Chemical Safety. Retrieved from <http://www.inchem.org/documents/ehc/ehc/ehc150.htm>

World Health Organization. (1993). Assessment of sources of air, water, and land pollution. A guide to rapid source inventory techniques and their use in formulating environmental control strategies. In Economopoulos, A. *Part One: Rapid inventory techniques in environmental pollution*. Retrieved from [http://whqlibdoc.who.int/Hq/1993/WHO\\_PEP\\_GETNET\\_93.1-A.pdf](http://whqlibdoc.who.int/Hq/1993/WHO_PEP_GETNET_93.1-A.pdf)

World Health Organization. (1995a). Briggs, D., Corvalán, & Nurminen, M. (Eds.). *Linkage methods for environment and health analysis. General guidelines*. Geneva, Switzerland: Author.

World Health Organization. (1995b). Inorganic lead. *Environmental Health Criteria*, 165. Geneva, Switzerland: Author. Retrieved from <http://www.inchem.org/documents/ehc/ehc/ehc165.htm>

World Health Organization. (1999). *Environmental epidemiology: a textbook on study methods and public health applications*. WHO/SDE/OEH/99.7. Geneva, Switzerland: Author.

World Health Organization. (2000a). Human exposure assessment. *Environmental Health Criteria*, 214. Retrieved from <http://www.inchem.org/documents/ehc/ehc/ehc214.htm>

World Health Organization. (2000b). *Guidelines for air quality*. Geneva, Switzerland: Author. Retrieved from [http://www.who.int/water\\_sanitation\\_health/dwq/GDWQ2004web.pdf](http://www.who.int/water_sanitation_health/dwq/GDWQ2004web.pdf)

---

World Health Organization. (2004). Ostro, B., (Ed), *Outdoor air pollution. Assessing the environmental burden of disease*. Retrieved from [http://www.who.int/quantifying\\_ehimpacts/publications/ebd5/en/](http://www.who.int/quantifying_ehimpacts/publications/ebd5/en/). [Accessed 6 October 2014]

World Health Organization. (2006). *Air quality guidelines – global update 2005*. Retrieved from [http://www.who.int/phe/health\\_topics/outdoorair/outdoorair\\_aqg/en/](http://www.who.int/phe/health_topics/outdoorair/outdoorair_aqg/en/)

World Health Organization. (2007). Wismar, M., Blau, J., Ernst, K., & Figueras, J., (Eds.), *The effectiveness of health impact assessment: Scope and limitations of supporting decision-making in Europe*. Retrieved from [http://www.euro.who.int/\\_\\_data/assets/pdf\\_file/0003/98283/E90794.pdf](http://www.euro.who.int/__data/assets/pdf_file/0003/98283/E90794.pdf).

World Health Organization. (2009). *Country profile of environmental burden of disease: China*. [http://www.who.int/entity/quantifying\\_ehimpacts/national/countryprofile/china.pdf](http://www.who.int/entity/quantifying_ehimpacts/national/countryprofile/china.pdf)

World Health Organization. (2010). Exposure to benzene: A major public health concern. Geneva, Switzerland: Author. Retrieved from: <http://www.who.int/ipcs/features/benzene.pdf>.

World Health Organization. (2011). *Burden of disease associated with urban outdoor air for 2008*. Retrieved from [http://www.who.int/phe/health\\_topics/outdoorair/databases/burden\\_disease/en/](http://www.who.int/phe/health_topics/outdoorair/databases/burden_disease/en/)

World Health Organization. (2013a). *Review of evidence on health aspects of air pollution –REVIHAAP Project* (Technical Report). Copenhagen, Denmark: WHO Regional Office for Europe. Retrieved from <http://www.euro.who.int/en/health-topics/environment-and-health/air-quality/publications/2013/review-of-evidence-on-health-aspects-of-air-pollution-revihaap-project-final-technical-report>.

World Health Organization. (2013b). *Health risks of air pollution in Europe – HRAPIE project. Recommendations for concentration-response functions for cost-benefit analysis of particulate matter, ozone and nitrogen dioxide*. Retrieved from <http://www.euro.who.int/en/health-topics/environment-and-health/air-quality/publications/2013/health-risks-of-air-pollution-in-europe-hrapie-project.-new-emerging-risks-to-health-from-air-pollution-results-from-the-survey-of-experts>.

World Health Organization. (2013c). *Chronic obstructive pulmonary disease* (Factsheet No. 315). Retrieved from <http://www.who.int/mediacentre/factsheets/fs315/en/>; [http://www.who.int/topics/chronic\\_obstructive\\_pulmonary\\_disease/en/](http://www.who.int/topics/chronic_obstructive_pulmonary_disease/en/).

World Health Organization. (2014a). *Ambient (outdoor) air quality and health* (Factsheet No. 313). Retrieved from [www.who.int/mediacentre/factsheets/fs313/en](http://www.who.int/mediacentre/factsheets/fs313/en)

World Health Organization. (2014b). *Burden of disease from ambient air pollution for 2012*. Retrieved from [http://www.who.int/phe/health\\_topics/outdoorair/databases/AAP\\_BoD\\_results\\_March2014.pdf?ua=1](http://www.who.int/phe/health_topics/outdoorair/databases/AAP_BoD_results_March2014.pdf?ua=1)

World Health Organization. (2014c). *7 million premature deaths annually linked to air pollution (Press release)*. Retrieved from <http://www.who.int/mediacentre/news/releases/2014/air-pollution/en/>

World Health Organization. (2014d). *Public health surveillance*. Retrieved from [http://www.who.int/topics/public\\_health\\_surveillance/en/](http://www.who.int/topics/public_health_surveillance/en/)

World Health Organization. (2015). *Health impact assessment (HIA) – tools and methods*. Retrieved from <http://www.who.int/hia/about/guides/en/>

---

World Health Organization/International Agency for Research on Cancer. (2013a). *Outdoor air pollution a leading environmental cause of cancer deaths* (Press release no. 221). Retrieved from [http://www.iarc.fr/en/media-centre/iarcnews/pdf/pr221\\_E.pdf](http://www.iarc.fr/en/media-centre/iarcnews/pdf/pr221_E.pdf)

World Health Organization/International Agency for Research on Cancer . (2013b). Air pollution and cancer. In Straif, K., Cohen, A., Samet, J. (Eds.) *IARC Scientific Publication*, No. 161. Retrieved from <http://www.iarc.fr/en/publications/books/sp161/index.php>

World Health Organization/International Agency for Research on Cancer. (2013c). Outdoor air pollution. *IARC monographs on the evaluation of carcinogenic risks to humans*, Volume 109. Lyon, France: IARC.

Yang G.H., Wang, Y., Zeng, Y.X., et al. (2013). Rapid health transition in China, 1990–2010: findings from the Global Burden of Disease Study 2010. *Lancet*, 381: 1987–2015.



Integrated Programme for  
Better Air Quality in Asia

[www.cleanairasia.org/ibaq](http://www.cleanairasia.org/ibaq)

ISBN 978-621-95434-0-8



9 786219 543408